

A DUO TO TALK ABOUT

BOTOX[®] COSMETIC
onabotulinumtoxinA

JUVÉDERM[®]
COLLECTION OF FILLERS

Contact us to discuss the treatment journey that's right for you.

LEARN MORE

Whitney was treated with BOTOX[®] Cosmetic for her moderate to severe frown lines, crow's feet lines, and forehead lines, along with JUVÉDERM[®] VOLUMA[®] XC in her cheeks and chin. Results may vary. The safety and efficacy of these products for combined use have not been studied.

Please see BOTOX[®] Cosmetic [Important Safety Information](#), including Boxed Warning, below.

BOTOX[®] Cosmetic

Only BOTOX[®] Cosmetic is FDA approved to temporarily improve the look of 3 areas—moderate to severe frown lines, crow's feet lines, and forehead lines—in adults.

The Look of 3 is a BOTOX[®] Cosmetic treatment of 3 areas, 64 Units, at least 3 times a year, spaced out at least 90 days apart.

Let's see if it's right for you.

EXPLORE NOW

JUVÉDERM[®] Collection of Fillers

Only JUVÉDERM[®] offers 5 long-lasting fillers to meet your unique needs. Add volume to temporarily lift and contour your cheeks, smooth lines around your nose and mouth, and plump lips.

SEE HOW

See Approved Uses and Important Safety Information below.

CONTACT US TODAY

Book your consultation

FOLLOW ALLÉ



DOWNLOAD OUR APP



ALLÉ MEMBER SUPPORT

1.888.912.1572

BOTOX[®] Cosmetic (onabotulinumtoxinA) Important Safety Information and Approved Uses

IMPORTANT SAFETY INFORMATION

BOTOX[®] Cosmetic may cause serious side effects that can be life threatening. Get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX[®] Cosmetic:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are pre-existing before injection. Swallowing problems may last for several months.
- **Spread of toxin effects.** The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms including: loss of strength and all-over muscle weakness, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice, trouble saying words clearly, loss of bladder control, trouble breathing, and trouble swallowing.

BOTOX[®] Cosmetic dosing units are not the same as, or comparable to, any other botulinum toxin product.

There has not been a confirmed serious case of spread of toxin effect when BOTOX[®] Cosmetic has been used at the recommended dose to treat frown lines, crow's feet lines, and/or forehead lines.

BOTOX[®] Cosmetic may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of taking BOTOX[®] Cosmetic. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Serious and/or immediate allergic reactions have been reported. They include: itching, rash, red itchy welts, wheezing, asthma symptoms, or dizziness or feeling faint. Get medical help right away if you are wheezing or have asthma symptoms, or if you become dizzy or faint.

Do not receive BOTOX[®] Cosmetic if you: are allergic to any of the ingredients in BOTOX[®] Cosmetic (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as *Myobloc[®]* (rimabotulinumtoxinB), *Dysport[®]* (abobotulinumtoxinA), or *Xeomin[®]* (incobotulinumtoxinA); have a skin infection at the planned injection site.

Tell your doctor about all your muscle or nerve conditions, such as ALS or Lou Gehrig's disease, myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects including difficulty swallowing and difficulty breathing from typical doses of BOTOX[®] Cosmetic.

Tell your doctor about all your medical conditions, including: plans to have surgery; had surgery on your face; have trouble raising your eyebrows; drooping eyelids; any other abnormal facial change; are pregnant or plan to become pregnant (it is not known if BOTOX[®] Cosmetic can harm your unborn baby); are breast-feeding or plan to (it is not known if BOTOX[®] Cosmetic passes into breast milk).

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using BOTOX[®] Cosmetic with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX[®] Cosmetic in the past.**

Tell your doctor if you have received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin; such as *Myobloc[®]*, *Dysport[®]*, or *Xeomin[®]* in the past (tell your doctor exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine; take aspirin-like products or blood thinners.

Other side effects of BOTOX[®] Cosmetic include: dry mouth; discomfort or pain at the injection site; tiredness; headache; neck pain; and eye problems: double vision, blurred vision, decreased eyesight, drooping eyelids and eyebrows, swelling of your eyelids and dry eyes.

For more information refer to the Medication Guide or talk with your doctor.

To report a side effect, please call Allergan at 1-800-678-1605.

Approved Uses

BOTOX[®] Cosmetic is a prescription medicine that is injected into muscles and used to temporarily improve the look of moderate to severe forehead lines, crow's feet lines, and frown lines between the eyebrows in adults.

Please see BOTOX[®] Cosmetic full [Product Information](#) including [Boxed Warning and Medication Guide](#).

JUVÉDERM[®] Injectable Gel Fillers Important Information

APPROVED USES

JUVÉDERM[®] VOLUMA[®] XC injectable gel is for deep injection in the cheek area to correct age-related volume loss and for augmentation of the chin region to improve the chin profile in adults over 21.

JUVÉDERM[®] VOLLURE[®] XC, JUVÉDERM[®] Ultra Plus XC, and JUVÉDERM[®] Ultra XC injectable gels are for injection into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds. JUVÉDERM[®] VOLLURE[®] XC injectable gel is for adults over 21.

JUVÉDERM[®] VOLBELLA[®] XC injectable gel is for injection into the lips for lip augmentation and for correction of perioral lines in adults over 21.

JUVÉDERM[®] Ultra XC injectable gel is for injection into the lips and perioral area for lip augmentation in adults over 21.

IMPORTANT SAFETY INFORMATION

Are there any reasons why I should not receive any JUVÉDERM[®] formulation?

Do not use these products if you have a history of multiple severe allergies or severe allergic reactions (anaphylaxis), or if you are allergic to lidocaine or the Gram-positive bacterial proteins used in these products.

What precautions should my doctor advise me about?

- Minimize strenuous exercise and exposure to extensive sun or heat within the first 24 hours following treatment. Exposure to any of these may cause temporary redness, swelling, and/or itching at the injection site
- Tell your doctor if you are pregnant or breastfeeding. The safety of these products for use during pregnancy or while breastfeeding has not been studied
- The safety of JUVÉDERM[®] VOLUMA[®] XC has not been studied in patients under 35 years or over 65 years for cheek augmentation, or under 22 years and over 80 years for chin augmentation. The safety of JUVÉDERM[®] VOLLURE[®] XC and JUVÉDERM[®] VOLBELLA[®] XC has not been studied in patients under 22 years, and the safety of JUVÉDERM[®] Ultra Plus XC and JUVÉDERM[®] Ultra XC has not been studied in patients under 18 years
- JUVÉDERM[®] VOLUMA[®] XC is intended for use in the chin and cheek areas. JUVÉDERM[®] VOLLURE[®] XC, JUVÉDERM[®] Ultra Plus XC, and JUVÉDERM[®] Ultra XC are intended for use in facial wrinkles and folds. JUVÉDERM[®] VOLBELLA[®] XC and JUVÉDERM[®] Ultra XC are intended for use in the lips and perioral area. The safety and effectiveness for treatment in other areas have not been established in clinical studies
- Tell your doctor if you have a history of excessive scarring (thick, hard scars) or pigmentation disorders. The safety of JUVÉDERM[®] products has not been studied in these patients and may result in additional scars or changes in pigmentation
- Tell your doctor if you are on therapy used to decrease the body's immune response (immunosuppressive therapy). Use may result in an increased risk of infection
- Tell your doctor before treatment if you are using substances that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners. As with any injection, this may result in increased bruising or bleeding at the injection site
- Patients who experience skin injury near the site of injection may be at a higher risk for adverse events
- JUVÉDERM[®] VOLUMA[®] XC was not studied in patients with significant loose skin of the chin, neck, or jaw
- The effect of JUVÉDERM[®] VOLUMA[®] XC injection into the chin on facial hair growth has not been studied

What are possible side effects?

The most commonly reported side effects with JUVÉDERM[®] injectable gels included redness, swelling, pain, tenderness, firmness, lumps/bumps, bruising, discoloration, and itching. For JUVÉDERM[®] VOLBELLA[®] XC, dryness was also reported. For JUVÉDERM[®] VOLLURE[®] XC, JUVÉDERM[®] Ultra Plus XC, and JUVÉDERM[®] Ultra XC injectable gels, most resolved within 14 days or less. For JUVÉDERM[®] VOLBELLA[®] XC, most resolved within 30 days or less. These side effects are consistent with other facial injection procedures.

Most side effects will resolve with time. Your doctor may choose to treat side effects persisting over 30 days with antibiotics, steroids, or hyaluronidase (an enzyme that breaks down hyaluronic acid).

One of the risks with these products is unintentional injection into a blood vessel. The chances of this happening are very small, but if it does happen, the complications can be serious and may be permanent. These complications, which have been reported for facial injections, can include vision abnormalities, blindness, stroke, temporary scabs, or permanent scarring of the skin.

As with all skin injection procedures, there is a risk of infection.

Visit [Juvederm.com](#) or talk to your doctor for more information. To report a side effect with any JUVÉDERM[®] product, please call Allergan at 1-800-433-8871.

Products in the JUVÉDERM[®] Collection are available only by a licensed physician or properly licensed practitioner.

Allergan Aesthetics

an AbbVie company

© 2022 AbbVie. All rights reserved. Allē[®] is a registered service mark of Allergan, Inc., an AbbVie company. BOTOX[®] and its design are registered trademarks of Allergan, Inc., an AbbVie company. JUVÉDERM[®] and its design are trademarks of Allergan Holdings France SAS, an AbbVie company, or its affiliates. Allergan Aesthetics reserves the right to alter or cancel the program with no advance notice or obligation. 2525 Dupont Drive, Irvine, CA 92612. SOLICITATION. MBD152774 01/22

Please do not reply to this email. If you would like to contact Allergan Aesthetics, please mail all communications to: Allergan, P.O. Box 19534, Irvine, CA 92623, USA.

Allē

EMAIL

If you no longer wish to receive emails from this sender, please reply to this email and include "unsubscribe" in the subject line.

LEGAL

[Privacy Statement](#) | [California Privacy Policy](#) | [AllerganAesthetics.com](#)